

Integral Homology of PGL_2 over Elliptic Curves

Kevin P. Knudson

The Friedlander–Milnor Conjecture [1] asserts that if G is a reductive algebraic group over an algebraically closed field k , then the comparison map

$$H_{\text{ét}}^{\bullet}(BG_k, \mathbb{Z}/p) \longrightarrow H^{\bullet}(BG, \mathbb{Z}/p)$$

is an isomorphism for all primes p not equal to the characteristic of k . Gabber’s rigidity theorem [2] implies that this map is indeed an isomorphism for the stable general linear group GL (this is due to Suslin [6] for $k = \mathbb{C}$ and to Jardine [3] for arbitrary k). Similarly, a proof of an unstable version of rigidity would lead to a proof of the unstable Friedlander–Milnor Conjecture.

In this note we consider unstable rigidity for the group PGL_2 over an elliptic curve E . We assume that E is defined by the equation $F(x, y) = 0$, where

$$F(x, y) = y^2 + a_1xy + a_3y - x^3 - a_2x^2 - a_4x - a_6,$$

and the a_i lie in an infinite field k . Denote by $\overline{E}$ the projective curve $E \cup \{\infty\}$. Denote by A the coordinate ring of the affine curve E . If $l \in k$ and $F(l, y) = 0$ has no rational solutions, denote by $k(\omega)$ the quadratic extension of k inside the algebraic closure $\overline{k}$ for which $F(l, \omega) = 0$. Our main result is the following.


THEOREM. *For all $i \geq 1$,*

$$\begin{aligned} H_i(PGL_2(A), \mathbb{Z}) &= \bigoplus_{\substack{p \in \overline{E} \\ 2p=0}} H_i(PGL_2(k), \mathbb{Z}) \oplus \bigoplus_{\substack{p \in \overline{E}, 2p \neq 0 \\ p \sim -p}} H_i(k^{\times}, \mathbb{Z}) \\ &\oplus \bigoplus_{\substack{l \in k, F(l, y)=0 \\ \text{has no rat. sol.}}} H_i(k(\omega)^{\times}/k^{\times}, \mathbb{Z}). \end{aligned}$$

Here, $p \sim -p$ means that we identify the factors corresponding to the points p and $-p$.

1991 *Mathematics Subject Classification.* 20G10.

Supported by an NSF Postdoctoral Fellowship, grant no. DMS–9627503.

FIGURE 1. $F(l, y) = 0$ has no rational solutionsFIGURE 2. $F(l, y) = 0$ has a unique rational solution

This is proved by examining the action of $PGL_2(A)$ on the Bruhat–Tits tree $\mathcal{X}$ associated to a two-dimensional vector space over the function field of E (see, *e.g.*, Serre’s book [5]). The quotient graph $PGL_2(A) \backslash \mathcal{X}$ is a tree (this is due to S. Takahashi [8]) and the various simplex stabilizers are easily described.

As a corollary, we have the following rigidity result.

COROLLARY. *Suppose the field k is algebraically closed and let x, y be distinct points on $\overline{E}$. Then the specialization homomorphisms*

$$s_x, s_y : H_\bullet(PGL_2(A), \mathbb{Z}) \longrightarrow H_\bullet(PGL_2(k), \mathbb{Z})$$

coincide.

1. The Quotient and the Stabilizers

In [8], Takahashi described a fundamental domain for the $PGL_2(A)$ -action on $\mathcal{X}$; denote this subtree by $\mathcal{D}$. There is a distinguished vertex $o \in \mathcal{D}$. For each l in $k \cup \{\infty\}$ there is a vertex $v(l)$ adjacent to o . The rest of $\mathcal{D}$ may be described as follows. Let $\mathcal{D}(l)$ denote the subtree of $\mathcal{D} - \{o\}$ which contains $v(l)$. The tree $\mathcal{D}$ is the union of o and the various $\mathcal{D}(l)$ (which are disjoint). The trees $\mathcal{D}(l)$ are as follows.

(1) Suppose $F(x, y) = 0$ has no rational solution with $x = l$. Then $\mathcal{D}(l)$ consists only of $v(l)$ (see Figure 1).


(2) Suppose $l = \infty$ or $F(x, y) = 0$ has a unique rational solution with $x = l$. Let p be the point at infinity of E or the rational point corresponding to the solution. Note that p is a point of order 2. Then $\mathcal{D}(l)$ consists of an infinite path $c(p, 1), c(p, 2), \dots$ and an extra vertex $e(p)$ (see Figure 2).

(3) Suppose $F(x, y) = 0$ has two different solutions such that $x = l$. Let p, q be the corresponding points on E . Then $\mathcal{D}(l)$ consists of two infinite paths $c(p, 1), c(p, 2), \dots$ and $c(q, 1), c(q, 2), \dots$ (see Figure 3).

The infinite path $c(p, 1), c(p, 2), \dots$ is called a *cusp*. Note that there is a one-to-one correspondence between cusps and the rational points of $\overline{E}$.

Since $\mathcal{X}$ is contractible, we have a spectral sequence with E^1 -term

$$E_{p,q}^1 = \bigoplus_{\sigma^{(p)} \subset \mathcal{D}} H_q(\Gamma_\sigma, \mathbb{Z}) \implies H_{p+q}(PGL_2(A), \mathbb{Z})$$

FIGURE 3. $F(l, y) = 0$ has two distinct solutions

where Γ_σ is the stabilizer of the p -simplex σ in $PGL_2(A)$.

The various stabilizers of the $GL_2(A)$ action were described in [8]. Denote these by $\tilde{\Gamma}_\sigma$.

PROPOSITION 1.1 (cf. [8], Theorem 5). *The stabilizers $\tilde{\Gamma}_\sigma$ are (up to isomorphism)*

$$\begin{aligned} \tilde{\Gamma}_o &\cong k^\times \\ \tilde{\Gamma}_{v(l)} &\cong \begin{cases} k(\omega)^\times & \text{in case (1)} \\ k^\times \times k & \text{in case (2)} \\ k^\times \times k^\times & \text{in case (3)} \end{cases} \\ \tilde{\Gamma}_{e(p)} &\cong GL_2(k) \\ \tilde{\Gamma}_{c(p,n)} &\cong (k^n \times_\theta k^\times) \times k^\times \end{aligned}$$

where $k^n \times_\theta k^\times$ is the semidirect product of k^n and $k^\times$ and θ is the automorphism of k^n given by coordinate-wise multiplication by elements of $k^\times$. Furthermore, the stabilizer of an edge is the intersection of the stabilizers of its vertices.

The groups $\tilde{\Gamma}_{c(p,n)}$ are of the form

$$\left\{ \begin{pmatrix} p & q \\ 0 & s \end{pmatrix} : p, s \in k^\times, q \in k^n \right\}.$$

Denote the diagonal subgroups of these $\tilde{\Gamma}$ by $\tilde{L}$. By Theorem 1.11 of [4], these stabilizers satisfy

$$H_{\bullet}(\tilde{\Gamma}, \mathbb{Z}) \cong H_{\bullet}(\tilde{L}, \mathbb{Z}),$$

the isomorphism being induced by the inclusion $\tilde{L} \rightarrow \tilde{\Gamma}$.

COROLLARY 1.2. *The stabilizers Γ_{σ} satisfy*

$$\begin{aligned} H_{\bullet}(\Gamma_o, \mathbb{Z}) &\cong H_{\bullet}(\{1\}, \mathbb{Z}) \\ H_{\bullet}(\Gamma_{v(l)}, \mathbb{Z}) &\cong \begin{cases} H_{\bullet}(k(\omega)^{\times}/k^{\times}, \mathbb{Z}) & \text{in case (1)} \\ H_{\bullet}(k, \mathbb{Z}) & \text{in case (2)} \\ H_{\bullet}(k^{\times}, \mathbb{Z}) & \text{in case (3)} \end{cases} \\ H_{\bullet}(\Gamma_{c(p,n)}, \mathbb{Z}) &\cong H_{\bullet}(k^{\times}, \mathbb{Z}) \\ H_{\bullet}(\Gamma_{e(p)}, \mathbb{Z}) &\cong H_{\bullet}(PGL_2(k), \mathbb{Z}). \end{aligned}$$

2. The Main Theorem

Note that our spectral sequence consists of two columns and that each row $E_{\bullet,q}^1$ is the chain complex $C_{\bullet}(\mathcal{D}, \mathcal{H}_q)$, where $\mathcal{H}_q$ is the coefficient system $\sigma \mapsto H_q(\Gamma_{\sigma})$. Fix a positive integer $q \geq 1$. Note that $H_q(\Gamma_o, \mathbb{Z}) = 0$ and that $H_q(\Gamma_e, \mathbb{Z}) = 0$ for any edge incident with o . It follows that the chain complex $C_{\bullet}(\mathcal{D}, \mathcal{H}_q)$ is a direct sum of chain complexes

$$C_{\bullet}(\mathcal{D}, \mathcal{H}_q) = \bigoplus_{l \in k \cup \{\infty\}} C_{\bullet}(\mathcal{D}(l), \mathcal{H}_q).$$

PROPOSITION 2.1. *Suppose $F(x, y) = 0$ has no rational solutions with $x = l$. Then*

$$H_i(\mathcal{D}(l), \mathcal{H}_q) = \begin{cases} H_q(k(\omega)^{\times}/k^{\times}, \mathbb{Z}) & i = 0 \\ 0 & i > 0. \end{cases}$$

PROOF. This is clear since $C_{\bullet}(\mathcal{D}(l), \mathcal{H}_q)$ consists only of the single group $H_q(k(\omega)^{\times}/k^{\times}, \mathbb{Z})$ sitting in degree zero. $\square$

PROPOSITION 2.2. *Suppose $l = \infty$ or $F(x, y) = 0$ has a unique rational solution with $x = l$. Then*

$$H_i(\mathcal{D}(l), \mathcal{H}_q) = \begin{cases} H_q(PGL_2(k), \mathbb{Z}) & i = 0 \\ 0 & i > 0. \end{cases}$$

PROOF. The stabilizer of $v(l)$ and of the edge joining $v(l)$ to $c(p, 1)$ is isomorphic to k . The map $\Gamma_{\langle v(l), c(p, 1) \rangle} \rightarrow \Gamma_{v(l)}$ is an isomorphism on homology and the map $\Gamma_{\langle v(l), c(p, 1) \rangle} \rightarrow \Gamma_{c(p, 1)}$ induces the zero map on homology. It follows that $H_{\bullet}(\mathcal{D}(l), \mathcal{H}_q) \cong H_{\bullet}(\mathcal{D}(l)', \mathcal{H}_q)$ where $\mathcal{D}(l)'$ is the tree obtained by deleting $v(l)$ and the edge joining it to $c(p, 1)$. One checks easily that the map

$$C_1(\mathcal{D}(l), \mathcal{H}_q) \longrightarrow C_0(\mathcal{D}(l), \mathcal{H}_q)$$

is injective with cokernel $H_q(PGL_2(k), \mathbb{Z})$ (or equivalently, check that the relative homology groups $H_\bullet(\mathcal{D}(l), e(p); \mathcal{H}_q)$ vanish). $\square$

PROPOSITION 2.3. *Suppose $F(x, y) = 0$ has two distinct solutions with $x = l$. Then*

$$H_i(\mathcal{D}(l), \mathcal{H}_q) = \begin{cases} H_q(k^\times, \mathbb{Z}) & i = 0 \\ 0 & i > 0. \end{cases}$$

PROOF. In this case, $\mathcal{D}(l)$ is a tree and the stabilizer of each vertex and each edge is $k^\times$. The maps $\Gamma_e \rightarrow \Gamma_v$ induce isomorphisms on homology for each edge and vertex. It follows that $C_\bullet(\mathcal{D}(l), \mathcal{H}_q)$ is a chain complex with *constant* coefficients. Since $\mathcal{D}(l)$ is contractible, the result follows. $\square$

THEOREM 2.4. *For all $i \geq 1$,*

$$\begin{aligned} H_i(PGL_2(A), \mathbb{Z}) \cong & \bigoplus_{\substack{p \in \overline{E} \\ 2p=0}} H_i(PGL_2(k), \mathbb{Z}) \oplus \bigoplus_{\substack{p \in \overline{E}, 2p \neq 0 \\ p \sim q}} H_i(k^\times, \mathbb{Z}) \\ & \oplus \bigoplus_{\substack{l \in k, F(l, y)=0 \\ \text{has no solutions}}} H_i(k(\omega)^\times / k^\times, \mathbb{Z}). \end{aligned}$$

PROOF. Note that since $\mathcal{D}$ is contractible, $E_{0,0}^2 = \mathbb{Z}$ and $E_{1,0}^2 = 0$. The preceding propositions show that

$$H_0(\mathcal{D}, \mathcal{H}_q) = \bigoplus_{l \in k \cup \{\infty\}} H_0(\mathcal{D}(l), \mathcal{H}_q)$$

and $H_1(\mathcal{D}, \mathcal{H}_q) = 0$. It remains to identify the various direct summands with points of $\overline{E}$.

Those l for which $F(l, y) = 0$ has a unique solution (or $l = \infty$) correspond to points of order 2 in $\overline{E}$. Those l for which $F(l, y) = 0$ has two distinct solutions correspond to pairs of points on $\overline{E}$. For such a pair p, q , the groups $H_i(k^\times, \mathbb{Z})$ arising from the cusps associated to p, q are identified together since they are both adjacent to $v(l)$. The claimed direct sum decomposition follows. $\square$

3. Rigidity

Suppose now that the field k is algebraically closed. Note that the projective curve $\overline{E}$ is isomorphic to the group $\text{Pic}^0 \overline{E}$ of degree zero line bundles on $\overline{E}$. Moreover, after a suitable linear change of coordinates, the points p, q corresponding to the two solutions of $F(l, y) = 0$ may be assumed to satisfy $q = -p$.

COROLLARY 3.1. *If k is algebraically closed, then for all $i \geq 1$,*

$$H_i(PGL_2(A), \mathbb{Z}) \cong \bigoplus_{\substack{L \in \text{Pic}^0 \overline{E} \\ 2L=0}} H_i(PGL_2(k), \mathbb{Z}) \oplus \bigoplus_{\substack{L \in \text{Pic}^0 \overline{E}, 2L \neq 0 \\ L \sim -L}} H_i(k^\times, \mathbb{Z}).$$

PROOF. This is obvious once one notes that the factors $H_i(k(\omega)^\times/k^\times, \mathbb{Z})$ do not enter the picture when k is algebraically closed. $\square$

As a consequence, we have the following rigidity result.

COROLLARY 3.2. *Suppose k is algebraically closed and let x, y be distinct points on $\overline{E}$. Then the corresponding specialization homomorphisms*

$$s_x, s_y : H_i(PGL_2(A), \mathbb{Z}) \longrightarrow H_i(PGL_2(k), \mathbb{Z})$$

coincide for all $i \geq 0$.

PROOF. This follows from the direct sum decomposition of Corollary 3.1. Since the groups which appear as summands do not involve rational functions on $\overline{E}$, the homomorphisms s_x and s_y must agree on each summand. $\square$

ACKNOWLEDGMENTS

I thank Andrei Suslin for pointing out that rigidity follows from the main theorem. I also thank Rick Jardine for many interesting discussions about the Friedlander–Milnor Conjecture.

References

- [1] E. Friedlander, G. Mislin, *Cohomology of classifying spaces of complex Lie groups and related discrete groups*, Comment. Math. Helv. **49** (1984), 347–361.
- [2] O. Gabber, *K-theory of Henselian local rings and Henselian pairs*, Contemp. Math. **126** (1992), 59–70.
- [3] J. Jardine, *Simplicial objects in a Grothendieck topos*, Contemp. Math. **55** (1986), 193–239.
- [4] Yu. Nesterenko, A. Suslin, *Homology of the full linear group over a local ring, and Milnor’s K-theory*, Math. USSR Izvestiya **34** (1990), 121–145.
- [5] J.-P. Serre, *Trees*, Springer-Verlag, Berlin/ Heidelberg, New York, 1980.
- [6] A. Suslin, *On the K-theory of local fields*, J. Pure Appl. Algebra **34** (1984), 301–318.
- [7] A. Suslin, *Algebraic K-theory of fields*, Proc. ICM, Berkeley, 1986.
- [8] S. Takahashi, *The fundamental domain of the tree of $GL(2)$ over the function field of an elliptic curve*, Duke Math. J. **73** (1993), 85–97.

DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY, EVANSTON, IL 60208
E-mail address: knudson@math.nwu.edu